

Comment on “Kerr Black Holes as Particle Accelerators to Arbitrarily High Energy”

Emanuele Berti^{1,2}, Vitor Cardoso^{2,3}, Leonardo Gualtieri⁴, Frans Pretorius⁵, Ulrich Sperhake¹

¹ *California Institute of Technology, Pasadena, CA 91109, USA*

² *Department of Physics and Astronomy, The University of Mississippi, University, MS 38677, USA*

³ *CENTRA, Departamento. de Física, Instituto. Superior. Técnico, Av. Rovisco Pais 1, 1049 Lisboa, Portugal*

⁴ *Dipartimento di Fisica, Università di Roma “Sapienza” & Sezione INFN Roma1, P.A. Moro 5, 00185, Roma, Italy and*

⁵ *Department of Physics, Princeton University, Princeton, NJ 08544, USA*

It has been suggested that rotating black holes could serve as particle colliders with arbitrarily high center-of-mass energy. Astrophysical limitations on the maximal spin, back-reaction effects and sensitivity to the initial conditions impose severe limits on the likelihood of such collisions.

PACS numbers: 97.60.Lf, 04.70.-s

Bañados, Silk and West (henceforth BSW) [1] recently observed that collisions of point particles falling from rest into rotating (Kerr) black holes (BHs) may have arbitrarily large center-of-mass (CM) energies close to the event horizon if the BH is maximally spinning and one of the particles has orbital angular momentum close to the value $L = L_{\text{scat}}$ corresponding to marginally bound geodesics.

An important practical limitation on the achievable CM energies occurs because, as pointed out by Thorne [2], the dimensionless spin of astrophysical BHs should not exceed $a/M = 0.998$. For $a/M = 0.998$, Eq. (14) in BSW yields maximum CM energies of about 10μ for particles of rest mass μ .

Even in the idealized scenario of *being given* an extremal BH, back-reaction effects make high-energy scattering very unlikely. Neglecting gravitational radiation, upon absorption of a pair of colliding particles of mass μ the dimensionless spin must be reduced by $\epsilon \equiv 1 - a/M \sim \mu/M$. After this first collision, Eq. (14) in BSW predicts that the new maximum allowed CM energy would be $E_{\text{CM}} \lesssim 10^{12} (\mu/1\text{MeV})^{3/4} (M/100M_{\odot})^{1/4}$ GeV, orders of magnitude below the Planck scale for typical values of the parameters. The collision of (say) a *single* electron pair would reduce the spin of a $100M_{\odot}$ BH enough to inhibit any further Planck-scale collisions. A hypothetical dark matter particle would need a mass $\mu \gtrsim 10^3$ TeV to allow for more than one Planck-scale event.

The estimates in BSW neglect gravitational radiation, which will significantly affect geodesics with $\delta = 1 - L/L_{\text{scat}} \ll 1$ (notice that in the *critical* case $L = L_{\text{scat}}$ it takes an infinite amount of proper time for a particle to reach the horizon). For small δ , the particle orbits around the marginally bound circular geodesic of a Schwarzschild BH $N \simeq -(\sqrt{2}\pi)^{-1} \log \delta$ times. For near-extremal Kerr BHs we get $N \sim -(2\pi\sqrt{2}\epsilon)^{-1} \log(8\sqrt{\epsilon}\delta)$. This simple analysis suggests that for $\delta \ll 1$ the radiation should be peaked at frequencies corresponding to marginally bound quasi-circular orbits with orbital frequency Ω_{mb} and that the total radiated energy $E_{\text{tot}} \sim N \sim -\log \delta$.

In Fig. 1 we confirm these conclusions by a numerical calculation of the energy radiated by particles of mass $\mu \ll M$ plunging from rest into a Schwarzschild BH (see [3] for details and notation). The dominant (quadrupo-




FIG. 1: Energy spectrum for $l = m = 2$ and different values of L/L_{scat} (as indicated in the legend). Inset: for small δ , a fit of the numerics yields $\mu^{-2} M E_{\text{tot}} \sim -0.11 - 0.18 \log \delta$.

lar) mode of the radiation shows a “bump,” as expected, at $\omega = 2\Omega_{\text{mb}} = (4M)^{-1}$. Since $E_{\text{tot}} \sim -\log \delta$ as $\delta \rightarrow 0$, radiative effects cannot be neglected in the analysis.

In conclusion, frame dragging effects in Kerr BHs can *in principle* accelerate particle to high energies, but (1) astrophysical restrictions on the spin severely limit the maximum CM energies in the collisions; (2) just as radiative losses constrain the performance of particle accelerators, gravitational radiation and back-reaction constrain the maximum CM energy for collisions around Kerr BHs; (3) the exponential sensitivity of whirling orbits to initial conditions requires significant fine-tuning to get sensible cross sections for the highest-energy collisions.

Acknowledgements. We thank M. Bañados, J. Silk and S. West for useful comments. This work was partially supported by NSF grants PHY-0900735, PHY-0745779, PHY-0601459, PHY-0652995, PHY-090003, FCT PTDC/FIS/64175/2006, PTDC/FIS/098025/2008, PTDC/FIS/098032/2008, the Alfred P. Sloan Foundation and the Sherman Fairchild Foundation.

-
- [1] M. Banados, J. Silk and S. M. West, Phys. Rev. Lett. **103**, 111102 (2009); arXiv:0909.0169 [hep-ph].
- [2] K. S. Thorne, Astrophys. J. **191**, 507 (1974).
- [3] K. I. Oohara and T. Nakamura, Prog. Theor. Phys. **70**, 757 (1983).